

Fig. 1

Clone 1 cDNA: malate dehydrogenase 2  
Clone ID: 1500012M15 3

DNA sequence

..... 4

Clone 2 cDNA: isocitrate dehydrogenase (NAD)  
Clone ID: 1500012E04

DNA sequence

.....

Clone 3 cDNA: isocitrate dehydrogenase (NADP)  
Clone ID: E030024J03

DNA sequence

.....

Clone 4 cDNA: oxoglutarate dehydrogenase (lipoamine)  
Clone ID: E430020N12

DNA sequence

.....

#### Procedures

To amplify cDNA inserts, cut out the 4 mm X 4 mm area spotted with DNA, place it into a PCR tube, add 25  $\mu$ l of water, centrifuge the resulting solution, and then initiate PCR cycle. The spot contains Taq DNA polymerase, an aptamer against the Taq DNA polymerase, trehalose, a PCR buffer, PCR primers (-21M13: 5'-TGTAAACGACGGCCAGT-3', 1233-Rv:5'-AGCGGATAACAATTTCACACAGGA-3'), each of dATP, dGTP, dCTP and dTTP, and MgCl<sub>2</sub>. After centrifuging the resulting solution, the PCR cycle is initiated. PCR cycles comprise 2 min at 94°C, 29 cycles of denaturing (94°C, 1 min), annealing (60°C, 1 min) and extension (68°C, 75 sec), and 15 min at 74°C.




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9





Fig. 10



Fwd T7 loxP Sfi I Sal I \*\*\*\*\* Bam HI SfI T3 Rev  
TGTAAAACGACGCCAGTGAATTGTAATACGACTCACTATAGGGCGAATTGGAGC  
TCCACCGCGGTGGCCCCCCCCCATAACTTCGTATAGCATACATTATACGAAGTTATG  
GATCAGGCCAATTCGGCCGAGCTCGAATTTCGTCGAC\*\*\*\*\*  
\*\*\*\*\*  
CGATCCGGCCATAAGGGCCTGATCCTTCGAGGGGGCCCGTACCAAGCTTTGT  
TGAGTTTACGAGGGTTAATTTCGAGCTTGGCGTAATCAGGTACAGCTGTTTC  
CTGRTGAAATTGTT

Fig. 11

cDNA: malate dehydrogenase  
Clone ID: 1500012M15

DNA sequence

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

Amino acid sequence

.....  
.....  
.....  
.....  
.....


Primer

-21M13 : 5'-TGTAAAACGACGGCCAGT-3'  
1233-Rv : 5'-AGCGGATAACAATTTCACACAGGA-3'

malate dehydrogenase cDNA  
primers



Fig. 12



BEST AVAILABLE COPY

Fig. 13

Amplification for human luteinizing hormone

## DNA sequence

**Primer**  
 – 21M13 : 5'-TGTAAACGACGCCAGT-3'  
 1233-Rv : 5'-AGCGGATAAACATTACACAGGA-3'

## Amplification mixture for human luteinizing hormone

Fig. 14

- Lane 1:primer set 1, aptamer (-)
- Lane 2:primer set 2, aptamer (-)
- Lane 3:size marker
- Lane 4:primer set 1, aptamer(+)
- Lane 5:primer set 2, aptamer(+)




Fig. 15

Clone 1 cDNA: malate dehydrogenase  
Clone ID: 1500012M15

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 2 cDNA: isocitrate dehydrogenase (NAD)  
Clone ID: 1500012E04

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 3 cDNA: isocitrate dehydrogenase (NADP)  
Clone ID: E030024J03

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 4 cDNA: oxoglutarate dehydrogenase (lipoamine)  
Clone ID: E430020N12

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 1      
Clone 2      
Clone 3      
Clone 4    

Chris Mizumoto, Reg. No. 42,899, 650-849-6684  
Atty. Docket #10084-0002-00000 Customer # 22,852  
For: "SUPPORT, PRINTED MATERIAL AND REAGENT...  
15/15

Fig. 16





Fig. 17



BEST AVAILABLE COPY